



Lassale, C., Tzoulaki, I., Moons, K. G. M., Sweeting, M., Boer, J., Johnson, L., & Butterworth, A. S. (2018). Separate and combined associations of obesity and metabolic health with coronary heart disease: a pan-European case-cohort analysis. *European Heart Journal*, 39(5), 397–406. [ehx448].
<https://doi.org/10.1093/eurheartj/ehx448>

Peer reviewed version

Link to published version (if available):
[10.1093/eurheartj/ehx448](https://doi.org/10.1093/eurheartj/ehx448)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via OUP at <https://academic.oup.com/eurheartj/article/4081012/Separate-and-combined-associations-of-obesity-and?searchresult=1#94987132>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Supplementary Material

Table of contents

Tables

p.2 – Table S1. Baseline characteristics and metabolic health status in members of the subcohort, by country

p.3 – Table S2. Sensitivity analysis: HR for CHD in different complete-case samples specific to each analysis.

p.4 – Table S3. Sensitivity analysis: HR for CHD where missing values are imputed by multiple imputation

p.5 – Table S4. Sensitivity analysis: HR for CHD after exclusion of first two years of follow-up across metabolically-defined body size phenotypes

p.6 – Table S5. Sensitivity analysis: HR for hard CHD (myocardial infarction) across metabolically-defined body size phenotypes

p.7 – Table S6. Sensitivity analysis: HR for CHD across metabolically-defined body size phenotypes in non-smokers only

p.8 – Table S7. Sensitivity analysis: HR for CHD events with the highest level of certainty across metabolically-defined body size phenotypes

p.9 – Table S8. Sensitivity analysis: HR for CHD across metabolically-defined body size phenotypes separately for men and women

p.10 – Table S9. Sensitivity analysis: HR for CHD in metabolically-defined body size phenotypes where the definition of MetS does not include the waist circumference criterion

p.11 – Table S10. Sensitivity analysis: HR for CHD in metabolically-defined body size phenotypes where “metabolically healthy” is defined as having none of the 4 abnormalities

p.12 – Table S11. Sensitivity analysis: HR for CHD in metabolically-defined body size phenotypes where obesity is defined by WC, and MetS does not include criteria on WC

p.13 – Table S12. Cross-classification in metabolically-defined body size phenotypes where body size is defined by BMI or by WC

Figures

p.14 – Figure S1. Schematic representation of the EPIC-CVD case-cohort design and sample included in the complete-case analysis

p.15 – Figure S2. Schematic representation of the analysis strategy

p.16 – Figure S3. Country-specific HRs across metabolically-defined body size phenotypes compared to metabolically healthy normal weight, Model B

Table S1. Baseline characteristics ^a and metabolic health status in members of the subcohort, by country

	Denmark	Greece	Germany	Italy	Netherlands	Spain	Sweden	United Kingdom
N	1895	1124	1356	1800	1270	550	1481	998
Age (years)	56.6 (4.4)	52.2 (12.2)	50 (8.7)	50.3 (7.9)	52.7 (10.7)	50.6 (8.4)	57.6 (7.7)	57 (10.7)
Women (%)	46.7	61.7	60.5	66.3	83.9	68.0	61.7	60.6
MetS (%)	27.3	28.1	29.3	22.1	24.1	26.6	25.7	22.1
Normal weight (%)	43.8	27.9	45.6	45.7	51.8	23.3	51.7	51.5
Overweight (%)	42.6	42.0	39.3	39.4	36.4	46.9	37.0	37.5
Obese (%)	13.7	30.1	15.1	14.8	11.8	29.8	11.3	11.0
MHO (% of the obese)	38.6	55.0	31.7	46.8	48.7	57.9	37.1	40.9

^a Values are unadjusted means (SD) or percentages.

n= 10,474 members of the subcohort included in the analytical sample

Abbreviations: MHO, metabolically healthy obese; MetS, metabolic syndrome

Table S2. Sensitivity analysis: HR for CHD in different complete-case samples specific to each analysis.

		HR ^a	95% CI	p	N cases	N total	I ² ^b	95% CI
BMI and Waist Circumference								
Model 1 ^c								
BMI		1.24	(1.17, 1.30)	<.0001	9212	23634	55%	(6%, 79%)
Waist circumference		1.31	(1.24, 1.38)	<.0001	9212	23634	47%	(0%, 76%)
Model 2 ^d								
BMI		1.05	(0.97, 1.14)	0.23	9212	23634	33%	(0%, 69%)
Waist circumference		1.24	(1.12, 1.37)	<.0001	9212	23634	40%	(0%, 73%)
Model 3 ^e								
BMI		1.06	(1.01, 1.10)	0.01	8319	18700	6%	(0%, 68%)
Waist circumference		1.07	(1.01, 1.14)	0.01	8319	18700	21%	(0%, 63%)
Metabolically-defined body size phenotypes								
Model B ^f								
BMI	MetS							
Normal weight	Metabolically Healthy	1.00	(ref)		2833	7664		
Overweight	Metabolically Healthy	1.26	(1.15, 1.38)	<.0001	2513	6748	0%	(0%, 68%)
Obese	Metabolically Healthy	1.41	(1.15, 1.72)	<.0001	687	2159	47%	(0%, 77%)
Normal weight	Metabolically Unhealthy	1.98	(1.67, 2.35)	<.0001	530	929	0%	(0%, 68%)
Overweight	Metabolically Unhealthy	2.26	(1.90, 2.67)	<.0001	2172	3703	63%	(19%, 83%)
Obese	Metabolically Unhealthy	2.44	(2.11, 2.82)	<.0001	1356	2495	4%	(0%, 69%)

^a Country-specific HRs were estimated from Prentice-weighted Cox proportional hazards models, and 95%CI estimated with robust variance, to take into account the case-cohort design. HRs were combined by multivariate random-effect meta-analysis across 8 countries. Age was used as the underlying time scale, models were stratified by sex and centre.

^b Heterogeneity across 8 European countries.

^c Model 1. HRs adjusted for age, smoking, physical activity, Mediterranean diet score, energy and alcohol intake, educational level

^d Model 2. Model 1 + waist circumference (for BMI) or BMI (for waist circumference)

^e Model 3. HRs adjusted for age, smoking, systolic blood pressure, total cholesterol, HDL cholesterol, history of diabetes

^f Model B. HRs adjusted for age, smoking, educational level, physical activity, Mediterranean diet score, energy and alcohol intake

Table S3. Sensitivity analysis: HR for CHD where missing values are imputed by multiple imputation

		HR ^a	95% CI	p
BMI and Waist Circumference				
Model 1 ^b				
BMI		1.28	(1.24, 1.33)	<.0001
Waist circumference		1.32	(1.27, 1.39)	<.0001
Model 2 ^c				
BMI		1.16	(1.07, 1.24)	<.0001
Waist circumference		1.15	(1.05, 1.25)	0.002
Model 3 ^d				
BMI		1.11	(1.06, 1.16)	<.0001
Waist circumference		1.10	(1.04, 1.16)	<.0001
Metabolically-defined body size phenotypes				
Model B ^e				
BMI	MetS			
Normal weight	Metabolically Healthy	1.00	(ref)	
Overweight	Metabolically Healthy	1.35	(1.20, 1.52)	<.0001
Obese	Metabolically Healthy	1.67	(1.39, 1.99)	<.0001
Normal weight	Metabolically Unhealthy	1.78	(1.46, 2.18)	<.0001
Overweight	Metabolically Unhealthy	2.22	(1.98, 2.49)	<.0001
Obese	Metabolically Unhealthy	2.43	(2.09, 2.81)	<.0001

^a HRs were estimated from Prentice-weighted Cox proportional hazards models, and 95%CI estimated with robust variance, to take into account the case-cohort design. Age was used as the underlying time scale, models were stratified by sex and centre. N=25,653 (12,240 cases). 5 imputed datasets, results combined by Rubin's rules.

^b Model 1. HRs adjusted for age, smoking, physical activity, Mediterranean diet score, energy and alcohol intake, educational level

^c Model 2. Model 1 + waist circumference (for BMI) or BMI (for waist circumference)

^d Model 3. HRs adjusted for age, smoking, systolic blood pressure, total cholesterol, HDL cholesterol, history of diabetes

^e Model B. HRs adjusted for age, smoking, physical activity, Mediterranean diet score, energy and alcohol intake, educational level

Table S4. Sensitivity analysis: HR for CHD after exclusion of first two years of follow-up across metabolically-defined body size phenotypes

BMI	MetS	HR ^a	95%CI	p-value	N cases	N total	I² ^b	95% CI
Model A ^c								
Normal weight	Metabolically Healthy	1.00	(ref)		1802	5961		
Overweight	Metabolically Healthy	1.22	(1.10, 1.35)	<.0001	1581	4241	0%	(0%, 68%)
Obese	Metabolically Healthy	1.24	(0.99, 1.57)	0.07	323	1059	37%	(0%, 72%)
Normal weight	Metabolically Unhealthy	2.12	(1.74, 2.58)	<.0001	428	778	0%	(0%, 68%)
Overweight	Metabolically Unhealthy	2.30	(1.96, 2.72)	<.0001	1678	2946	50%	(0%, 78%)
Obese	Metabolically Unhealthy	2.54	(2.23, 2.91)	<.0001	976	1829	0%	(0%, 68%)
Model B ^d								
Normal weight	Metabolically Healthy	1.00	(ref)		1802	5961		
Overweight	Metabolically Healthy	1.23	(1.11, 1.37)	<.0001	1581	4241	0%	(0%, 68%)
Obese	Metabolically Healthy	1.25	(0.99, 1.58)	0.06	323	1059	35%	(0%, 71%)
Normal weight	Metabolically Unhealthy	2.09	(1.72, 2.54)	<.0001	428	778	0%	(0%, 68%)
Overweight	Metabolically Unhealthy	2.29	(1.90, 2.75)	<.0001	1678	2946	58%	(7%, 81%)
Obese	Metabolically Unhealthy	2.46	(2.14, 2.82)	<.0001	976	1829	0%	(0%, 68%)

^a Country-specific HRs were estimated from Prentice-weighted Cox proportional hazards models, and 95%CI estimated with robust variance, to take into account the case-cohort design. HRs were combined by multivariate random-effects meta-analysis across 8 countries. Age was used as the underlying time scale, models were stratified by sex and centre. n=16,814 (6,788 CHD cases).

^b Heterogeneity across 8 European countries

^c Model A. HRs adjusted for age, smoking, educational level.

^d Model B included the same variables as model A + physical activity, Mediterranean diet score, energy and alcohol intake

Table S5. Sensitivity analysis: HR for hard CHD (myocardial infarction) across metabolically-defined body size phenotypes

BMI	MetS	HR ^a	95%CI	p-value	N cases	N total	I² ^b	95% CI
Model A ^c								
Normal weight	Metabolically Healthy	1.00	(ref)		1303	6165		
Overweight	Metabolically Healthy	1.23	(1.07, 1.41)	0.003	1144	4451	0%	(0%, 68%)
Obese	Metabolically Healthy	1.32	(1.01, 1.73)	0.045	234	1103	43%	(0%, 75%)
Normal weight	Metabolically Unhealthy	2.13	(1.70, 2.67)	<.0001	325	842	12%	(0%, 71%)
Overweight	Metabolically Unhealthy	2.21	(1.92, 2.55)	<.0001	1283	3200	19%	(0%, 62%)
Obese	Metabolically Unhealthy	2.57	(2.17, 3.04)	<.0001	773	1972	12%	(0%, 72%)
Model B ^d								
Normal weight	Metabolically Healthy	1.00	(ref)		1303	6165		
Overweight	Metabolically Healthy	1.24	(1.08, 1.43)	0.002	1144	4451	3%	(0%, 69%)
Obese	Metabolically Healthy	1.30	(1.00, 1.70)	0.049	234	1103	39%	(0%, 73%)
Normal weight	Metabolically Unhealthy	2.10	(1.71, 2.58)	<.0001	325	842	0%	(0%, 68%)
Overweight	Metabolically Unhealthy	2.21	(1.89, 2.57)	<.0001	1283	3200	26%	(0%, 67%)
Obese	Metabolically Unhealthy	2.46	(2.10, 2.90)	<.0001	773	1972	1%	(0%, 68%)

^a Country-specific HRs were estimated from Prentice-weighted Cox proportional hazards models, and 95%CI estimated with robust variance, to take into account the case-cohort design. HRs were combined by multivariate random-effects meta-analysis across 8 countries. Age was used as the underlying time scale, models were stratified by sex and centre. n=17,733 participants (5,062 CHD cases)

^b Heterogeneity across 8 European countries

^c Model A. HRs adjusted for age, smoking, educational level.

^d Model B included the same variables as model A + physical activity, Mediterranean diet score, energy and alcohol intake

Table S6. Sensitivity analysis: HR for CHD events across metabolically-defined body size phenotypes in non-smokers only

BMI	MetS	HR ^a	95%CI	p-value	N cases	N total	I² ^b	95% CI
Model A ^c								
Normal weight	Metabolically Healthy	1.00	(ref)		1193	4166		
Overweight	Metabolically Healthy	1.22	(1.04, 1.38)	0.003	1173	3215	0%	(0%,68%)
Obese	Metabolically Healthy	1.26	(0.96, 1.67)	0.10	272	871	45%	(0%,76%)
Normal weight	Metabolically Unhealthy	2.15	(1.72, 2.69)	<.0001	277	506	0%	(0%,68%)
Overweight	Metabolically Unhealthy	2.32	(2.00, 2.70)	<.0001	1233	2193	7%	(0%,70%)
Obese	Metabolically Unhealthy	2.59	(2.21, 3.03)	<.0001	763	1443	0%	(0%,68%)
Model B ^d								
Normal weight	Metabolically Healthy	1.00	(ref)		1193	4166		
Overweight	Metabolically Healthy	1.23	(1.08,1.40)	0.002	1173	3215	47%	(0%, 80%)
Obese	Metabolically Healthy	1.29	(0.99,1.67)	0.06	272	871	0%	(0%, 79%)
Normal weight	Metabolically Unhealthy	2.15	(1.71,2.71)	<.0001	277	506	35%	(0%, 75%)
Overweight	Metabolically Unhealthy	2.32	(1.97,2.73)	<.0001	1233	2193	44%	(0%, 80%)
Obese	Metabolically Unhealthy	2.58	(2.18,3.04)	<.0001	763	1443	0%	(0%, 79%)

^a Country-specific HRs were estimated from Prentice-weighted Cox proportional hazards models, and 95%CI estimated with robust variance, to take into account the case-cohort design. HRs were combined by multivariate random-effects meta-analysis across 8 countries. Age was used as the underlying time scale, models were stratified by sex and centre. n=12,394 (4,911 cases)

^b Heterogeneity across 8 European countries

^c Model A. HRs age, smoking (never, former), educational level.

^d Model B included the same variables as model A + physical activity, Mediterranean diet score, energy and alcohol intake

Table S7. Sensitivity analysis: HR for CHD events with the highest level of certainty across metabolically-defined body size phenotypes

BMI	MetS	HR ^a	95%CI	p-value	N cases	N total	I ² ^b	95% CI
Model A ^c								
Normal weight	Metabolically Healthy	1.00	(ref)		268	3159		
Overweight	Metabolically Healthy	1.39	(1.04, 1.86)	0.03	325	2598	40%	(0%, 78%)
Obese	Metabolically Healthy	1.11	(0.79, 1.55)	0.55	65	738	0%	(0%, 79%)
Normal weight	Metabolically Unhealthy	2.58	(1.68, 3.95)	<.0001	75	374	21%	(0%, 66%)
Overweight	Metabolically Unhealthy	2.78	(2.17, 3.57)	<.0001	389	1707	3%	(0%, 80%)
Obese	Metabolically Unhealthy	2.99	(2.36, 3.79)	<.0001	262	1218	0%	(0%, 79%)
Model B ^d								
Normal weight	Metabolically Healthy	1.00	(ref)		268	3159		
Overweight	Metabolically Healthy	1.44	(1.05, 1.99)	0.03	325	2598	47%	(0%, 80%)
Obese	Metabolically Healthy	1.13	(0.81, 1.59)	0.47	65	738	0%	(0%, 79%)
Normal weight	Metabolically Unhealthy	2.79	(1.73, 4.49)	<.0001	75	374	35%	(0%, 75%)
Overweight	Metabolically Unhealthy	2.88	(2.09, 3.98)	<.0001	389	1707	44%	(0%, 80%)
Obese	Metabolically Unhealthy	2.93	(2.29, 3.77)	<.0001	262	1218	0%	(0%, 79%)

^a Country-specific HRs were estimated from Prentice-weighted Cox proportional hazards models, and 95%CI estimated with robust variance, to take into account the case-cohort design. HRs were combined by multivariate random-effects meta-analysis across 8 countries. Age was used as the underlying time scale, models were stratified by sex and centre. n=9,794 (1,384 cases)

^b Heterogeneity across 8 European countries

^c Model A. HRs adjusted for age, smoking, educational level.

^d Model B included the same variables as model A + physical activity, Mediterranean diet score, energy and alcohol intake

Table S8. Sensitivity analysis: HR for CHD across metabolically-defined body size phenotypes separately for men and women

BMI	MetS	HR ^a	95%CI	p-value	N cases	N total	I ² ^b	95% CI
Men								
Normal weight	Metabolically Healthy	1.00	(ref)		1053	2310		
Overweight	Metabolically Healthy	1.32	(1.16, 1.51)	<.0001	1168	2328	0%	(0%,68%)
Obese	Metabolically Healthy	1.28	(0.99, 1.65)	0.06	184	415	7%	(0%,70%)
Normal weight	Metabolically Unhealthy	2.53	(1.90, 3.37)	<.0001	243	346	45%	(0%,76%)
Overweight	Metabolically Unhealthy	2.35	(2.03, 2.73)	<.0001	1239	1861	25%	(0%,66%)
Obese	Metabolically Unhealthy	2.47	(2.06, 2.95)	<.0001	622	954	0%	(0%,68%)
Women								
Normal weight	Metabolically Healthy	1.00	(ref)		925	3855		
Overweight	Metabolically Healthy	1.19	(1.03, 1.38)	0.02	609	2123	0%	(0%,68%)
Obese	Metabolically Healthy	1.28	(1.02, 1.60)	0.03	176	688	0%	(0%,68%)
Normal weight	Metabolically Unhealthy	1.85	(1.48, 2.31)	<.0001	248	496	0%	(0%,68%)
Overweight	Metabolically Unhealthy	2.36	(2.01, 2.76)	<.0001	677	1339	26%	(0%,67%)
Obese	Metabolically Unhealthy	2.72	(2.27, 3.26)	<.0001	493	1018	30%	(0%,69%)

^a Country-specific HRs were estimated from Prentice-weighted Cox proportional hazards models, and 95%CI estimated with robust variance, to take into account the case-cohort design. HRs were combined by multivariate random-effects meta-analysis across 8 countries. Age was used as the underlying time scale, models were stratified by sex and centre. n=8,214 men (4,509 cases) and n=9,519 (n=3,128 cases).

HRs adjusted for age, smoking (never, former), educational level, physical activity, Mediterranean diet score, energy and alcohol intake

^b Heterogeneity across 8 European countries

Table S9. Sensitivity analysis: HR for CHD in metabolically-defined body size phenotypes where the definition of MetS does not include the waist circumference criterion

BMI	MetS without WC	HR ^a	95%CI	p-value	N cases	N total	I ² ^b	95% CI
Model A ^c								
Normal weight	Metabolically Healthy	1.00	(ref)		1458	5166		
Overweight	Metabolically Healthy	1.36	(1.20, 1.56)	<.0001	1532	3999	6%	(0%, 69%)
Obese	Metabolically Healthy	1.43	(1.13, 1.81)	0.003	357	1095	42%	(0%, 74%)
Normal weight	Metabolically Unhealthy	2.01	(1.72, 2.35)	<.0001	1011	1841	19%	(0%, 62%)
Overweight	Metabolically Unhealthy	2.52	(2.17, 2.94)	<.0001	2161	3652	41%	(0%, 74%)
Obese	Metabolically Unhealthy	2.92	(2.54, 3.34)	<.0001	1118	1980	0%	(0%, 68%)
Model B ^d								
Normal weight	Metabolically Healthy	1.00	(ref)		1458	5166		
Overweight	Metabolically Healthy	1.38	(1.20, 1.57)	<.0001	1532	3999	7%	(0%, 70%)
Obese	Metabolically Healthy	1.43	(1.14, 1.81)	0.002	357	1095	39%	(0%, 73%)
Normal weight	Metabolically Unhealthy	2.00	(1.72, 2.32)	<.0001	1011	1841	8%	(0%, 70%)
Overweight	Metabolically Unhealthy	2.49	(2.11, 2.94)	<.0001	2161	3652	48%	(0%, 77%)
Obese	Metabolically Unhealthy	2.82	(2.45, 3.25)	<.0001	1118	1980	0%	(0%, 68%)

^a Country-specific HRs were estimated from Prentice-weighted Cox proportional hazards models, and 95%CI estimated with robust variance, to take into account the case-cohort design. HRs were combined by multivariate random-effect meta-analysis across 8 countries. Age was used as the underlying time scale, models were stratified by sex and centre. n=17,733 participants (7,637 CHD cases)

^b Heterogeneity across 8 European countries

^c Model A. HRs adjusted for age, smoking, educational level.

^d Model B included the same variables as model A + physical activity, Mediterranean diet score, energy and alcohol intake

Table S10. Sensitivity analysis: HR for CHD in metabolically-defined body size phenotypes where “metabolically healthy” is defined as having none of the 4 abnormalities

BMI	Healthy defined as having 0 abnormality	HR ^a	95%CI	p-value	N cases	N total	I ² ^b	95% CI
Model A ^c								
Normal weight	Metabolically Healthy	1.00	(ref)		390	2148		
Overweight	Metabolically Healthy	1.24	(0.99, 1.56)	0.06	260	1040	0%	(0%, 68%)
Obese	Metabolically Healthy	1.15	(0.72, 1.84)	0.55	36	198	0%	(0%, 68%)
Normal weight	Metabolically Unhealthy	1.93	(1.62, 2.31)	<.0001	2079	4859	0%	(0%, 68%)
Overweight	Metabolically Unhealthy	2.67	(2.29, 3.12)	<.0001	3433	6611	0%	(0%, 68%)
Obese	Metabolically Unhealthy	3.19	(2.64, 3.85)	<.0001	1439	2877	0%	(0%, 68%)
Model B ^d								
Normal weight	Metabolically healthy	1.00	(ref)		390	2148		
Overweight	Metabolically Healthy	1.24	(0.99, 1.56)	0.06	260	1040	0%	(0%, 68%)
Obese	Metabolically Healthy	1.21	(0.76, 1.92)	0.43	36	198	0%	(0%, 68%)
Normal weight	Metabolically Unhealthy	1.94	(1.62, 2.32)	<.0001	2079	4859	0%	(0%, 68%)
Overweight	Metabolically Unhealthy	2.68	(2.28, 3.14)	<.0001	3433	6611	0%	(0%, 68%)
Obese	Metabolically Unhealthy	3.12	(2.57, 3.80)	<.0001	1439	2877	2%	(0%, 68%)

^a Country-specific HRs were estimated from Prentice-weighted Cox proportional hazards models, and 95%CI estimated with robust variance, to take into account the case-cohort design. HRs were combined by multivariate random-effects meta-analysis across 8 countries. Age was used as the underlying time scale, models were stratified by sex and centre. n=17,733 participants (7,637 CHD cases)

^b Heterogeneity across 8 European countries

^c Model A. HRs adjusted for age, smoking, educational level.

^d Model B included the same variables as model A + physical activity, Mediterranean diet score, energy and alcohol intake

Table S11. Sensitivity analysis: HR for CHD in metabolically-defined body size phenotypes where obesity is defined by WC, and MetS does not include criteria on WC

WC	MetS without WC	HR ^a	95%CI	p-value	N cases	N total	I ² ^b	95% CI
Model A ^c								
Normal WC ^d	Metabolically Healthy	1.00	(ref)		1653	5707		
Overweight ^e	Metabolically Healthy	1.40	(1.15, 1.69)	0.001	1031	2729	55%	(0%, 80%)
Obese ^f	Metabolically Healthy	1.37	(1.11, 1.69)	0.003	663	1824	50%	(0%, 78%)
Normal WC ^d	Metabolically Unhealthy	1.96	(1.70, 2.25)	<.0001	1143	2075	0%	(0%, 68%)
Overweight ^e	Metabolically Unhealthy	2.43	(2.09, 2.82)	<.0001	1340	2292	20%	(0%, 62%)
Obese ^f	Metabolically Unhealthy	2.92	(2.47, 3.45)	<.0001	1807	3106	47%	(0%, 76%)
Model B ^g								
Normal WC ^d	Metabolically Healthy	1.00	(ref)		1653	5707		
Overweight ^e	Metabolically Healthy	1.41	(1.17, 1.70)	<.0001	1031	2729	53%	(0%, 79%)
Obese ^f	Metabolically Healthy	1.39	(1.12, 1.73)	0.003	663	1824	51%	(0%, 78%)
Normal WC ^d	Metabolically Unhealthy	1.94	(1.68, 2.24)	<.0001	1143	2075	0%	(0%, 68%)
Overweight ^e	Metabolically Unhealthy	2.43	(2.09, 2.84)	<.0001	1340	2292	24%	(0%, 65%)
Obese ^f	Metabolically Unhealthy	2.84	(2.38, 3.39)	<.0001	1807	3106	50%	(0%, 77%)
Model C ^h								
Normal WC ^d	Metabolically Healthy	1.00	(ref)		1653	5707		
Overweight ^e	Metabolically Healthy	1.33	(1.10, 1.60)	<.0001	1031	2729	44%	(0%, 75%)
Obese ^f	Metabolically Healthy	1.22	(0.99, 1.51)	0.06	663	1824	25%	(0%, 66%)
Normal WC ^d	Metabolically Unhealthy	1.93	(1.66, 2.24)	<.0001	1143	2075	0%	(0%, 68%)
Overweight ^e	Metabolically Unhealthy	2.26	(1.92, 2.66)	<.0001	1340	2292	21%	(0%, 63%)
Obese ^f	Metabolically Unhealthy	2.44	(1.99, 2.99)	<.0001	1807	3106	25%	(0%, 66%)

^a Country-specific HRs were estimated from Prentice-weighted Cox proportional hazards models, and 95%CI estimated with robust variance, to take into account the case-cohort design. HRs were combined by multivariate random-effect meta-analysis across 8 countries. Age was used as the underlying time scale, models were stratified by sex and EPIC study centre. n=17,733 participants (7,637 CHD cases); ^b Heterogeneity across 8 countries;

^c Model A. HRs adjusted for age, smoking, educational level.;

^d Normal WC: WC<94 for men, 80 for women; ^e Overweight: 94≤WC<102 for men, 80≤WC<88 for women; ^f Obese: WC≥102 for men, 88 for women

^g Model B included the same variables as model A + physical activity, Mediterranean diet score, energy and alcohol intake

^h Model C included the same variables as model B + BMI

Table S12. Cross-classification in metabolically-defined body size phenotypes where body size is defined by BMI or by WC

BMI-defined	WC-defined						Total
	MHANW	MUANW	MHAOW	MUAOW	MHAO	MUAO	
MHNW	4447	999	648	0	71	0	6165
MUNW	0	394	0	391	0	57	842
MHOW	1246	452	1905	0	848	0	4451
MUOW	0	218	0	1753	0	1229	3200
MHO	14	8	176	0	905	0	1103
MUO	0	4	0	148	0	1820	1972
Total	5707	2075	2729	2292	1824	3106	17733

Abbreviations: MHNW, metabolically healthy normal weight; MUNW, metabolically unhealthy normal weight; MHOW, metabolically healthy overweight; MUOW, metabolically unhealthy overweight; MHO, metabolically healthy obese; MUO, metabolically unhealthy obese; MHANW, metabolically healthy abdominally normal weight; MUANW, metabolically unhealthy abdominally normal weight; MHAOW, metabolically healthy abdominally overweight; MUAOW, metabolically unhealthy abdominally overweight; MHAO, metabolically healthy abdominally obese; MUAO, metabolically unhealthy abdominally obese.

Weighted kappa (95%CI), measuring agreement between two classifications, was 0.667 (0.660- 0.674)

Figure S1. Schematic representation of the EPIC-CVD case-cohort design and sample included in the complete-case analysis




Figure S2. Schematic representation of the analysis strategy





Figure S3. Country-specific HRs across metabolically-defined body size phenotypes compared to metabolically healthy normal weight, Model B^a



^a Model B was adjusted for age, smoking, educational level, physical activity, Mediterranean diet score, energy and alcohol intake